

## Flavonoids and Triterpenes from the Nest of the Stingless Bee *Trigona spinipes*

Marinalva O. Freitas, Flávio A. F. Ponte, Mary Anne S. Lima and Edilberto R. Silveira\*

Curso de Pós-Graduação em Química Orgânica, Departamento de Química Orgânica e Inorgânica, Centro de Ciências, Universidade Federal do Ceará, CP 12 200, 60451-970 Fortaleza-CE, Brazil




Figure S1. <sup>1</sup>H NMR spectrum (500 MHz, MeOD) of magniferolic acid (1) and 3β-hydroxy-24-methylenecycloartan-26-oic acid (2)

\*e-mail: edil@ufc.br




Figure S2.  $^{13}\text{C}$ -BB NMR spectrum (125 MHz, MeOD) of magniferolic acid (1) and 3 $\beta$ -hydroxy-24-methylenecycloartan-26-oic acid (2)








Figure S3.  $^{13}\text{C}$ -DEPT 135° NMR spectrum (125 MHz, MeOD) of magniferolic acid (1) and 3 $\beta$ -hydroxy-24-methylenecycloartan-26-oic acid (2)



**Figure S4.** <sup>1</sup>H NMR spectrum (500 MHz, C<sub>5</sub>H<sub>5</sub>N) of 3'-methyl-quercetin (3)



**Figure S5.** <sup>13</sup>C-BB NMR spectrum (125 MHz, C<sub>5</sub>H<sub>5</sub>N) of 3'-methyl-quercetin (3)

Figure S6. <sup>1</sup>H NMR spectrum (500 MHz, MeOD) of sakuranetin (4)Figure S7. <sup>13</sup>C NMR spectrum (500 MHz, MeOD) of sakuranetin (4)




Figure S8. <sup>1</sup>H NMR spectrum (500 MHz, C<sub>3</sub>H<sub>5</sub>N) of kaempferol 7-methyl ether (5)






Figure S9. <sup>13</sup>C-BB NMR spectrum (125 MHz, C<sub>3</sub>H<sub>5</sub>N) of kaempferol 7-methyl ether (5)

Figure S10. <sup>1</sup>H NMR spectrum (500 MHz, MeOD) of tricetin (6)Figure S11. <sup>13</sup>C-BB NMR spectrum (500 MHz, MeOD) of tricetin (6)




Figure S12. <sup>1</sup>H NMR spectrum (500 MHz, MeOD) of aromadendrin 7-methyl ether (7)




Figure S13. <sup>13</sup>C -BB NMR spectrum (500 MHz, MeOD) of aromadendrin 7-methyl ether (7)