


## Supplementary Information

### Corrosion Behavior of NdFeB Magnets in Different Aqueous Solutions

Ana-Maria Popescu, <sup>a</sup> Jose M. Calderon-Moreno,<sup>a</sup> Kazimir Yanushkevich,<sup>b</sup>  
Alexei Aplevich,<sup>b</sup> Olga Demidenko, <sup>b</sup> Elena Ionela Neacsu<sup>a</sup> and Virgil Constantin <sup>\*,a</sup>

<sup>a</sup>“Ilie Murgulescu” Institute of Physical Chemistry of the Romanian Academy, Splaiul Independentei 202,  
Bucharest 060021, Romania

<sup>b</sup>SSPA “Scientific-Practical Materials Research Centre of NAS of Belarus”, 19 P. Brovki str.,  
Minsk 220072, Belarus



**Figure S1.** (a) Tetragonal unit cell of Nd<sub>2</sub>Fe<sub>14</sub>B (P42/mnm space group) and (b) trigonal prism containing a boron atom in the Nd<sub>2</sub>Fe<sub>14</sub>B structure (reproduced from reference 1 with copyright permission 2014 from John Wiley and Sons).

#### Reference

1. Jadhav, A. P.; Ma, H.; Kim, D. S.; Baek, Y. K.; Choi, C. J.; Kang, Y. S.; *Bull. Korean Chem. Soc.* **2014**, 35, 886. [Crossref]

\*e-mail: virgilconstantin@yahoo.co

