Supplementary Information

Generation of Volatile Compounds from Carotenoids of *Dunaliella bardawil* Algae by Water Bath Heating and Microwave Irradiation

Natália A. B. Tinoco,^a Thais M. Uekane,^a Anna Tsukui,^a Paula F. de Aguiar,^a Cláudia M. L. L. Teixeira^b and Claudia M. Rezende**^a

^aInstituto de Química, Universidade Federal do Rio de Janeiro, 21945-970 Rio de Janeiro-RJ, Brazil

^bInstituto Nacional de Tecnologia, 20081-312 Rio de Janeiro-RJ, Brazil




Figure S1. Chromatogram of volatile compounds present in *D. bardawil* biomass (BC2) obtained by GC-qMS. 1: β-cyclocitral; 2: α-ionone; 3: 7,8-dihydro-β-ionone; 4: *trans*-geranyl acetone; 5: 5,6-epoxy-β-ionone; 6: β-ionone; 7: dihydroactinidiolide.

^{*}e-mail: claudia.rezendeufrj@gmail.com




Figure S2. Response surface in absolute area obtained by GC-MS (SIM) for (a) α -ionone; (b) β -ionone; (c) β -cyclocitral as a function of temperature and time in a DVB/CAR/PDMS SPME fiber.

(b) (a) β-cyclocitral α-ionone y = 401.729,2477x + 681.224,54927,50E+06 2,50E+07 y = 133.361,2798x - 204.656,2921 $R^2 = 0,9832$ $R^2 = 0,9894$ 6,00E+06 2,00E+07 **4,**50E+06 3,00E+06 **5** 1,50E+07 1,00E+07 1,50E+06 5,00E+06 0,00E+00 0,00E+00 12,0 15,0 80 100 120 140 160 6,0 9,0 18,0 20 60 Concetration (ng mL -1) Concentration (ng mL-1) (c) (d) α-ionone β-cyclocitral 5,E+05 8,E+05 Residue 2,E+05 -1,E+05 3,E+05 -2,E+05 100 120 140 160 -4,E+05 -7,E+05

Figure S3. Calibration curves and residual plots for β -cyclocitral (a and c) and α -ionone (b and d), obtained by GC-qMS (SIM).

-1,E+06

Concentration (ng mL -1)

-7,E+05

Concentration (ng mL -1)




Figure S4. Typical chromatographic profile of *D. bardawil* biomass by HPLC-UV (460 nm). Lutein (1); chlorophyll b (2); chlorophyll a (3); β -carotene (4).




Figure S5. Response surface of the concentration of: (a) β -cyclocitral BU; (b) β -cyclocitral MW; (c) α -ionone BU; (d) α -ionone MW; (e) β -ionone BU; (f) β -ionone MW; (g) total carotenoids WB; (h) total carotenoids MW as a function of temperature and time employed for degradation of carotenoids from *D. bardawil*.

Table S1. Concentration (ng mL⁻¹) of the major volatile compounds produced on the conditions of the central points from MW and WB experiments

Volatile compound	Concentration / (ng mL ⁻¹)	
	MW (1.5 min)	WB (45 min)
β-Cyclocitral	4.46 ± 0.34	12.83 ± 0.11
α-Ionone	5.37 ± 0.85	15.27 ± 0.17
β-Ionone	67.56 ± 0.57	115.67 ± 0.63

MW: microwave irradiation; WB: water bath heating.